

UNIVERZITA MATEJA BELA V BANSKEJ BYSTRICI
FAKULTA PRÍRODNÝCH VIED
KATEDRA MATEMATIKY

O zväzoch kongruencií zväzov

(ŠVOČ)

AUTOR PRÁCE: DANIELA GUFFOVÁ

VEDÚCI PRÁCE: DOC. RNDR. MIROSLAV HAVIAR, CSc.

2009

UNIVERSITY OF MATEJ BEL IN BANSKÁ BYSTRICA
FACULTY OF NATURAL SCIENCES

DEPARTMENT OF MATHEMATICS

On Congruence Lattices of Lattices

(Student Competition)

AUTHOR: DANIELA GUFFOVÁ

SUPERVISOR: DOC. RNDR. MIROSLAV HAVIAR, CSc.

2009

I would like to thank my supervisor, Dr Miroslav Haviar, for giving me incentive ideas and facilitating my work.

CONTENTS

1. Introduction	1
2. Preliminaries	2
3. Lattices with relative Stone congruence lattices	3
4. Lattices with relative L_n -congruence lattices	5
5. Conclusion	11
References	12

1. INTRODUCTION

One of the basic facts about the congruence lattices of lattices is that they are distributive and pseudocomplemented. T. Tanaka [11], P. Crawley [1], G. Grätzer and E. T. Schmidt [3] have characterized those lattices whose congruence lattices are Boolean. In the monograph [2], G. Grätzer posed problems (problems III.5 and III.6) of characterizing those lattices whose congruence lattices considered as pseudocomplemented lattices belong to the n th Lee's equational class B_n of distributive pseudocomplemented lattices described by the identity

$$(L_n) \quad (x_1 \wedge \dots \wedge x_n)^* \vee (x_1^* \wedge \dots \wedge x_n)^* \vee \dots \vee (x_1 \wedge \dots \wedge x_n^*)^* = 1.$$

Distributive pseudocomplemented lattices satisfying the identity (L_n) are called (L_n) -lattices [6], [7]. As the class B_1 is the class of all Stone lattices, (L_1) -lattices are in fact Stone lattices. Lattices whose congruence lattices are Stone have been characterized by T. Katriňák [8]. Later, M. Haviar [6] characterized lattices with (L_n) -congruence lattices for arbitrary $n \geq 1$.

Distributive pseudocomplemented lattices in which every interval satisfies the identity (L_n) are called relative (L_n) -lattices. In [4], M. Haviar and T. Katriňák characterized lattices with relative Stone congruence lattices. Lattices with relative (L_n) -congruence lattices were characterized later by M. Haviar in [6]. Semi-discrete lattices with (L_n) - and relative (L_n) -congruence lattices were characterized by M. Haviar and T. Katriňák in [7].

The congruence lattices of lattices are also relatively pseudocomplemented, hence they can be investigated as Heyting algebras. It is natural to seek for a characterization of lattices whose congruence lattices satisfy identities formulated in terms of relative pseudocomplement. In particular, relative (L_n) -lattices can be characterized by the identity

$$(L'_n) \quad (x_1 \wedge \dots \wedge x_n) * y \vee (x_1 * y \wedge \dots \wedge x_n) * y \vee \dots \vee (x_1 \wedge \dots \wedge x_n * y) * y = 1.$$

In [7] only semi-discrete lattices whose congruence lattices satisfy the identity (L'_n) were described. In this work we present a description of arbitrary lattices whose congruence lattices considered as Heyting algebras satisfy the identity (L'_n) (section 4). In particular, one obtains a description of lattices with relative (L_1) -congruence lattices. In Section 3 we give a slightly different description of lattices with relative Stone congruence lattices than is the one obtained in Section 4 in case $n = 1$.

Our method is alternative to the one presented in [6] and [4] where the identity (L'_n) was not used and the respective descriptions of lattices with relative (L_n) - and relative Stone congruence lattices were presented by translating the corresponding conditions for factor lattices L/π (π is a congruence of L) with (L_n) - and Stone congruence lattices without the need to write down the proofs

for the given characterizations. In our approach presented here we entirely use the identities (L'_n) and we actually write down self-contained proofs for the characterizations of lattices with relative (L_n)- and relative Stone congruence lattices.

2. PRELIMINARIES

The following basic concepts and facts can be found in [2], [4], [7] or [6].

Let $\text{Con } L$ denote the lattice of all congruences on a lattice L with Δ and ∇ , the smallest and the largest congruence relation. The lattice $\text{Con } L$ is distributive, moreover $\text{Con } L$ satisfies the infinite distributivity law

$$\theta \wedge \bigvee (\alpha_i : i \in I) = \bigvee (\theta \wedge \alpha_i : i \in I)$$

for any $\theta, \alpha_i \in \text{Con } L$.

It follows that for any $\alpha, \beta \in \text{Con } L$ there exists a largest congruence δ such that $\alpha \wedge \delta \leq \beta$. It is obvious that $\delta = \bigvee (\sigma : \alpha \wedge \sigma \leq \beta)$. The congruence δ is called the *relative pseudocomplement of α with respect to β* and denoted by $\alpha * \beta$. Therefore $\langle \text{Con } L, \vee, \wedge, *, \Delta, \nabla \rangle$ is a complete relatively pseudocomplemented lattice, i.e. a complete Heyting algebra.

Recall that an algebra $\langle H, \vee, \wedge, *, 0, 1 \rangle$ of type $(2, 2, 2, 0, 0)$ is a *Heyting algebra* if it satisfies:

(H1) $\langle H, \vee, \wedge \rangle$ is a distributive lattice,

(H2) $x \wedge 0 = 0, x \vee 1 = 1,$

(H3) $x * x = 1,$

(H4) $(x * y) \wedge y = y, x \wedge (x * y) = x \wedge y,$

(H5) $x * (y \wedge z) = (x * y) \wedge (x * z), (x \vee y) * z = (x * z) \wedge (y * z).$

The Heyting algebras were introduced by G. Birkhoff under the name *Brouwerian algebras*.

K. B. Lee [10] has shown that the lattice of all equational subclasses of the class B_ω of all distributive pseudocomplemented lattices (*p-algebras*) is a chain

$$B_{-1} \subset B_0 \subset B_1 \subset \dots \subset B_n \subset \dots \subset B_\omega$$

of type $\omega + 1$, where B_{-1}, B_0, B_1 are the classes of all trivial p-algebras, Boolean algebras, and Stone algebras, respectively. Moreover, a distributive pseudocomplemented lattice belongs to the class B_n ($n \geq 1$) if and only if it satisfies the identity

$$(L_n) \quad (x_1 \wedge \dots \wedge x_n)^* \vee (x_1^* \wedge \dots \wedge x_n^*)^* \vee \dots \vee (x_1 \wedge \dots \wedge x_n^*)^* = 1,$$

i.e. is an (L_n)-lattice.

A distributive relatively pseudocomplemented lattice $\langle L, \vee, \wedge, *, 0, 1 \rangle$ is a relative Stone lattice if and only if

$$x * y \vee (x * y) * y = 1$$

for every $x, y \in L$. A distributive relatively pseudocomplemented lattice L is a relative (L_n) -lattice ($n \geq 1$) if and only if it satisfies the identity

$$(L'_n) \quad (x_1 \wedge \dots \wedge x_n) * y \vee (x_1 * y \wedge \dots \wedge x_n) * y \vee \dots \vee (x_1 \wedge \dots \wedge x_n * y) * y = 1.$$

One of the mostly used concepts in this work is the concept of weak projectivity of quotients. We denote a/b an ordered pair of elements a, b of a lattice L satisfying $b \leq a$; a/b is called a *quotient* of L . A quotient c/d is a *subquotient* of a/b if $b \leq d \leq c \leq a$. We call a/b a *proper quotient* if $b < a$. If $b \prec a$, i.e. b is covered by a , then a/b is called a *prime quotient*.

We will say that a quotient a/b is *weakly projective* to a quotient c/d and use the notation $a/b \rightarrow c/d$ if there exist finitely many elements $x_1, \dots, x_n \in L$ such that

$$\begin{aligned} c &= (\dots ((a \vee x_1) \wedge x_2) \vee \dots) \vee x_n, \\ d &= (\dots ((b \vee x_1) \wedge x_2) \vee \dots) \vee x_n. \end{aligned}$$

The importance of weak projectivity in the description of lattice congruences is given by the following two lemmas.

Lemma 2.1. ([6], Lemma 1) *For any principal congruence $\theta_{a,b} \in \text{Con } L$,*

$$(c, d) \in \theta_{a,b},$$

($d \leq c, b \leq a$) if and only if there is a finite chain $d = y_0 \leq \dots \leq y_n = c$ such that $a/b \rightarrow y_{i+1}/y_i$ for all $i \in \{0, \dots, n-1\}$.

Lemma 2.2. ([6], Lemma 2) *Let L be a lattice and $\theta, \varphi \in \text{Con } L$. then the relative pseudocomplement of θ with respect to φ is*

$$\theta * \varphi = \bigvee_{(\theta_{u,v}, (u,v) \in S)} (\theta_{u,v}, (u,v) \in S),$$

where S is the set of all pairs of elements (u, v) ($u, v \in L$) such that $u/v \rightarrow z/t$ and $(z, t) \in \theta$ implies $(z, t) \in \varphi$ for all $z, t \in L$.

3. LATTICES WITH RELATIVE STONE CONGRUENCE LATTICES

In this section we give a description of lattices with relative Stone congruence lattices.

Definition 3.1. ([4], Definition 1) *Let L be a lattice, $\pi \in \text{Con } L$ and $a/b, u/v$ quotients of L . Then L is said to be π -almost weakly modular whenever $a/b \rightarrow u/v$ and $(u, v) \notin \pi$ imply the existence of a subquotient $a_1/b_1 \subseteq a/b$ with $(a_1, b_1) \notin \pi$ such that for every quotient r/s with $a_1/b_1 \rightarrow r/s$ and $(r, s) \notin \pi$ there exists a quotient z/t with $r/s \rightarrow z/t$, $u/v \rightarrow z/t$ and $(z, t) \notin \pi$.*

Definition 3.2. ([4], Definition 2) *Let L be a lattice and $\theta, \pi \in \text{Con } L$, $\theta \geq \pi$. Then θ is said to be π -weakly separable if for any $a < b$ in L there exists a chain $a = z_0 \leq z_1 \leq \dots \leq z_n = b$ such that for each $i \in \{0, \dots, n-1\}$ either*

- (i) $z_{i+1}/z_i \rightarrow u/v$ and $(u, v) \in \theta$ imply $(u, v) \in \pi$ or
- (ii) for every subquotient $r/s \subseteq z_{i+1}/z_i$ with $(r, s) \notin \pi$, there exists a quotient u/v with $r/s \rightarrow u/v$ and $(u, v) \in \theta$, $(u, v) \notin \pi$.

Theorem 3.3. ([4], Theorem 2) *Let L be a lattice. The lattice $\text{Con } L$ is relative Stone if and only if for every $\pi \in \text{Con } L$ the following conditions hold:*

- (1) L is π -almost weakly modular and
- (2) every congruence $\theta \geq \pi$ is π -weakly separable.

Proof. First we will prove the necessity.

Let $\text{Con } L$ be relatively Stone lattice, i. e. it satisfies the identity

$$(\theta * \pi) \vee ((\theta * \pi) * \pi) = \nabla,$$

for all congruences $\theta, \pi \in \text{Con } L$. Let $a/b, u/v$ be quotients of L such that $a/b \rightarrow u/v$ with $(u, v) \notin \pi$ and $a > b, u > v$. Set

$$\phi := \theta_{u,v} \vee \pi.$$

Since $\text{Con } L$ is relatively Stone, it follows that

$$(a, b) \in (\phi * \pi) \vee ((\phi * \pi) * \pi) = (\theta_{u,v} * \pi) \vee ((\theta_{u,v} * \pi) * \pi),$$

so there exists a chain $b = c_0 \leq c_1 \leq \dots \leq c_n = a$ such that for every $i \in \{0, \dots, n-1\}$

$$(c_{i+1}, c_i) \in (\theta_{u,v} * \pi) \text{ or } (c_{i+1}, c_i) \in ((\theta_{u,v} * \pi) * \pi).$$

If for every $i \in \{0, \dots, n-1\}$ the first case holds, we get $(a, b) \in \theta_{u,v} * \pi$, that is, $(u, v) \in \theta_{u,v} * \pi$, so $(u, v) \in \pi$, a contradiction.

Thus there is a subquotient $a_1/b_1 \subseteq a/b$ such that $(a_1, b_1) \notin (\theta_{u,v} * \pi)$ and $(a_1, b_1) \in ((\theta_{u,v} * \pi) * \pi)$. Let r/s be a quotient such that $a_1/b_1 \rightarrow r/s$ and $(r, s) \notin \pi$. Then $(r, s) \in ((\theta_{u,v} * \pi) * \pi)$.

Whenever the conditions $r/s \rightarrow z'/t'$, $(z', t') \in \theta_{u,v}$ would imply $(z', t') \in \pi$ we would get $(r, s) \in (\theta_{u,v} * \pi)$, so $(r, s) \in \pi$ would also hold, a contradiction.

Hence there exists a quotient z'/t' such that $r/s \rightarrow z'/t'$, with $(z', t') \in \theta_{u,v}$ and $(z', t') \notin \pi$. Since $(z', t') \in \theta_{u,v}$, there exists a subquotient $z/t \subseteq z'/t'$ such that $u/v \rightarrow z/t$. As also $r/s \rightarrow z/t$, $\text{Con } L$ is π -almost weakly modular.

Now let $\theta \in \text{Con } L$ with $\theta \geq \pi$. Since $\text{Con } L$ is relative Stone lattice, $(a, b) \in (\theta * \pi) \vee ((\theta * \pi) * \pi)$ for any $a > b$. Therefore there exists a chain $b = z_0 \leq \dots \leq z_m = a$ such that

$$(z_{i+1}, z_i) \in (\theta * \pi) \text{ or } (z_{i+1}, z_i) \in ((\theta * \pi) * \pi).$$

In the first case we get that $z_{i+1}/z_i \rightarrow u/v$ and $(u, v) \in \theta$ implies $(u, v) \in \pi$. So we get the condition (i) from the Definition 3.2. Now

let $(z_{i+1}, z_i) \in ((\theta * \pi) * \pi)$. Let r/s be a subquotient of the quotient z_{i+1}/z_i with $(r, s) \notin \pi$ and let $r/s \rightarrow u/v$, $(u, v) \notin \pi$. If for every $u' \geq v'$ the conditions $u/v \rightarrow u'/v'$ and $(u', v') \in \theta$ imply $(u', v') \in \pi$, then $(u, v) \in (\theta * \pi)$. So we would get $(u, v) \in ((\theta * \pi) * \pi)$ and $(u, v) \in (\theta * \pi)$, which yields $(u, v) \in \pi$, a contradiction.

So there exists a quotient u'/v' such that $u/v \rightarrow u'/v'$, $(u', v') \in \theta$ and $(u', v') \notin \pi$. The π -weakly separability of any congruence $\theta \in \text{Con } L$ has been proved.

Now we will prove the sufficiency. Let $a < b$ and let $\theta, \pi \in \text{Con } L$, $\theta \geq \pi$. From π -weakly separability of θ follows the existence of a chain $a = z_0 \leq \dots \leq z_n = b$ such that for every $i \in \{0, \dots, n-1\}$ (i) or (ii) from Definition 3.2 holds. If (i) holds, we get $(z_i, z_{i+1}) \in (\theta * \pi)$. Now let assume that (i) from the Definition 3.2 does not hold and that (ii) from the Definition 3.2 holds for (z_i, z_{i+1}) . We will distinguish two cases:

I. Let assume that $z_i/z_{i+1} \rightarrow u/v$, $(u, v) \in (\theta * \pi)$ imply $(u, v) \in \pi$. We get $(z_i, z_{i+1}) \in ((\theta * \pi) * \pi)$.

II. There remains the case when $z_i/z_{i+1} \rightarrow u/v$, $(u, v) \in (\theta * \pi)$ but $(u, v) \notin \pi$. The π -almost weakly modularity of $\text{Con } L$ yields the existence of a subquotient $a_1/b_1 \subseteq z_i/z_{i+1}$, with $(a_1, b_1) \notin \pi$, such that for every quotient r/s and $a_1/b_1 \rightarrow r/s$ with $(r, s) \notin \pi$ there exists a quotient z/t with $u/v \rightarrow z/t$ and $r/s \rightarrow z/t$ with $(z, t) \notin \pi$. From (ii) of Definition 3.2 it follows that there exists a quotient u/v such that $a_1/b_1 \rightarrow u/v$, $(u, v) \in \theta$ and $(u, v) \notin \pi$. By π -almost weakly modularity of L there exists a quotient z/t such that $u/v \rightarrow z/t$ and $(z, t) \notin \pi$. Then $(z, t) \in \theta$ and $(z, t) \in \theta * \pi$, so $(z, t) \in \pi$, a contradiction. Therefore the case II. cannot occur, so for every $i \in \{0, \dots, n-1\}$

$$(z_{i+1}, z_i) \in (\theta * \pi) \text{ or } (z_{i+1}, z_i) \in ((\theta * \pi) * \pi)$$

holds. Hence $\text{Con } L$ is relative Stone lattice. \square

Theorem 3.3 yields the following statement.

Corollary 3.4. ([4], Theorem 1) *Let L be a lattice. Then $\text{Con } L$ is a Stone lattice if and only if the following conditions hold:*

- (1) *L is Δ -almost weakly modular and*
- (2) *every congruence of L is Δ -weakly separable.*

4. LATTICES WITH RELATIVE L_n -CONGRUENCE LATTICES

In this section we give we a description of arbitrary lattices whose congruence lattices considered as Heyting algebras satisfy the identity (L'_n) .

Definition 4.1. ([5], Definition 3) Let L be a lattice, $n \geq 1$ and $a/b, u_1/v_1, \dots, u_{n+1}/v_{n+1}$ be nontrivial quotients of L . Then L is said to be $(\pi - n)$ -**weakly modular** whenever

$$a/b \rightarrow u_i/v_i \text{ and } (u_i, v_i) \notin \pi, \quad i = 1, \dots, n+1$$

imply that one of the following conditions holds:

- (i) there exist $i, j \in \{1, \dots, n+1\}, i \neq j$ and a quotient u/v such that $u_i/v_i \rightarrow u/v, u_j/v_j \rightarrow u/v$ with $(u, v) \notin \pi$.
- (ii) for all $i \in \{1, \dots, n+1\}$ there is a proper subquotient $r_i/s_i \subset a/b$ such that $(r_i, s_i) \notin \pi$ and $(r_i, a) \notin \pi$ or $(s_i, b) \notin \pi$ and a quotient z_i/t_i , such that $r_i/s_i \rightarrow z_i/t_i, u_i/v_i \rightarrow z_i/t_i$ and $(z_i, t_i) \notin \pi$.

Definition 4.2. ([5], Definition 4) Let L be a lattice, $\pi \in \text{Con } L$ and $n \geq 1$. Then an (unordered) n -tuple $\theta_1, \dots, \theta_n$ ($\theta_1, \dots, \theta_n \geq \pi$) is said to be $(\pi - n)$ -**separable** if for any $b < a$ there exists a chain $b = z_0 \leq z_1 \leq \dots \leq z_n = a$ such that for every $i \in \{0, \dots, n-1\}$ either

- (i) $z_{i+1}/z_i \rightarrow u/v$ and $(u, v) \in (\theta_1 \cap \dots \cap \theta_n)$ imply $(u, v) \in \pi$ or
- (ii) there exists some $j \in \{1, \dots, n\}$ such that for every proper subquotient $r/s \subset z_{i+1}/z_i$ with $(r, s) \notin \pi$ and $(r, z_{i+1}) \notin \pi$ or $(s, z_i) \notin \pi$ the following holds: $(u, v) \in (\theta_1 \cap \dots \cap \theta_{j-1} \cap \theta_{j+1} \cap \dots \cap \theta_n)$, $r/s \rightarrow u/v$ and $(u, v) \notin \pi$ imply the existence of a quotient u'/v' such that $u/v \rightarrow u'/v'$ and $(u', v') \in \theta_j, (u', v') \notin \pi$.

Theorem 4.3. ([5], Theorem 4) Let L be a lattice and $n \geq 1$. $\text{Con } L$ is relative (L_n) -lattice if and only if for every $\pi \in \text{Con } L$ the following conditions hold:

- (i) L is $(\pi - n)$ -weakly modular and
- (ii) every n -tuple of congruences $\theta_1, \dots, \theta_n$ on L such that $\theta_i \geq \pi$ for all $i = 1, \dots, n$ is $(\pi - n)$ -separable.

Proof. Assume that $\text{Con } L$ satisfies the identity

$$(L'_n) \quad ((\theta_1 \wedge \dots \wedge \theta_n) * \pi) \vee ((\theta_1 * \pi \wedge \dots \wedge \theta_n) * \pi) \vee \dots \vee ((\theta_1 \wedge \dots \wedge \theta_n * \pi) * \pi) = \nabla.$$

We shall prove that L is $(\pi - n)$ -weakly modular. Let $\pi \in \text{Con } L$ and let $a/b, u_1/v_1, \dots, u_{n+1}/v_{n+1}$ be nontrivial quotients in L such that $a/b \rightarrow u_i/v_i$ and $(u_i, v_i) \notin \pi$ for $i = 1, \dots, n+1$. Consider there are no $i, j \in \{1, \dots, n+1\}, i \neq j$ and a quotient $u/v, (u, v) \notin \pi$ such that $u_i/v_i \rightarrow u/v, u_j/v_j \rightarrow u/v$. Set

$$\phi_1 := \theta_{u_1, v_1} \vee \pi, \quad \dots, \quad \phi_{n+1} := \theta_{u_{n+1}, v_{n+1}} \vee \pi.$$

We shall prove that

$$(1) \quad (\phi_1 * \pi) \vee \dots \vee (\phi_{n+1} * \pi) = \nabla.$$

We will show that $\phi_i \cap \phi_j = \pi$ for all $i, j \in \{1, \dots, n+1\}, i \neq j$. It is obvious that $\pi \subseteq \phi_i \cap \phi_j$. To prove the equality suppose the existence of elements $u, v \in L, u > v$, such that $(u, v) \notin \pi$ and $(u, v) \in (\theta_i \cap \theta_j)$. By distributivity we get $\phi_i \cap \phi_j = (\theta_{u_i, v_i} \wedge \theta_{u_j, v_j}) \vee \pi$. Thus there exists a chain $v = c_0 \leq \dots \leq c_n = u$ such that $(c_{k+1}, c_k) \in (\theta_{u_i, v_i} \wedge \theta_{u_j, v_j})$ or $(c_{k+1}, c_k) \in \pi$. Since $(u, v) \notin \pi$ there exists a nontrivial subquotient $u'/v' \subseteq u/v$ such that $(u', v') \in (\theta_{u_i, v_i} \wedge \theta_{u_j, v_j})$ and $(u', v') \notin \pi$. By Lemma 1 there exists a nontrivial subquotient $u''/v'' \subseteq u'/v'$ such that $u_i/v_i \rightarrow u''/v'', u_j/v_j \rightarrow u''/v''$ and $(u'', v'') \notin \pi$, a contradiction. Hence $\phi_i \cap \phi_j = \pi$ and $\phi_i \leq \phi_j * \pi$ for all $i, j \in \{1, \dots, n+1\}, i \neq j$.

In the case $n = 1$ we have $(\phi_1 * \pi) \vee ((\phi_1 * \pi) * \pi) = \nabla$. Since $\phi_2 \leq \phi_1 * \pi$, we get $\phi_2 * \pi \geq (\phi_1 * \pi) * \pi$. So

$$\nabla = (\phi_1 * \pi) \vee ((\phi_1 * \pi) * \pi) \leq (\phi_1 * \pi) \vee (\phi_2 * \pi),$$

thus (1) holds. Now assume $n \geq 2$. Set

$$\begin{aligned} \alpha_1 &:= \phi_2 \vee \phi_3 \vee \dots \vee \phi_n \vee \phi_{n+1} \\ \alpha_2 &:= \phi_1 \vee \phi_3 \vee \dots \vee \phi_n \vee \phi_{n+1} \\ &\vdots \\ \alpha_n &:= \phi_1 \vee \phi_2 \vee \dots \vee \phi_{n-1} \vee \phi_{n+1}. \end{aligned}$$

We have

$$((\alpha_1 \wedge \dots \wedge \alpha_n) * \pi) \vee ((\alpha_1 * \pi \wedge \dots \wedge \alpha_n) * \pi) \vee \dots \vee ((\alpha_1 \wedge \dots \wedge \alpha_n * \pi) * \pi) = \nabla.$$

We will prove that

$$(2) \quad (\alpha_1 \wedge \dots \wedge \alpha_n) = \phi_{n+1}, (\alpha_1 * \pi \wedge \dots \wedge \alpha_n) = \phi_1, \dots, (\alpha_1 \wedge \dots \wedge \alpha_n * \pi) = \phi_n.$$

First we will show that

$$\alpha_1 \wedge \dots \wedge \alpha_n = \phi_{n+1}.$$

Clearly $\phi_{n+1} \subseteq \alpha_1 \wedge \dots \wedge \alpha_n$. Suppose on the contrary that there exist $u, v \in L, (u, v) \in (\alpha_1 \wedge \dots \wedge \alpha_n)$ and $(u, v) \notin \phi_{n+1}$. As $(u, v) \in \alpha_1$, there exists some $i \in \{2, \dots, n\}$ and a subquotient $u'/v' \subseteq u/v, (u', v') \notin \pi$ such that $(u', v') \in \phi_i$ and $(u', v') \notin \phi_{n+1}$. We also have $(u', v') \in \alpha_i$, so there exist $j \in \{1, \dots, n\} - \{i\}$ and a subquotient $u''/v'' \subseteq u'/v', (u'', v'') \notin \pi$ such that $(u'', v'') \in \phi_j$. Then $(u'', v'') \in (\phi_i \cap \phi_j)$ that contradicts $\phi_j \cap \phi_i = \pi$, for $i \neq j$. Therefore

$$\alpha_1 \wedge \dots \wedge \alpha_n = \phi_{n+1}.$$

Also

$$\alpha_i * \pi = (\phi_1 * \pi) \wedge \dots \wedge (\phi_{i-1} * \pi) \wedge (\phi_{i+1} * \pi) \wedge \dots \wedge (\phi_{n+1} * \pi).$$

Using the fact that $\phi_i \cap \phi_j = \pi$, for all $i \neq j$ and the distributivity law we get

$$\alpha_1 \wedge \dots \wedge (\alpha_i * \pi) \wedge \dots \wedge \alpha_n = ((\phi_1 * \pi) \wedge \dots \wedge (\phi_{i-1} * \pi) \wedge \phi_i \wedge (\phi_{i+1} * \pi) \wedge \dots \wedge (\phi_n * \pi)) \vee \pi.$$

As $\phi_i \leq \phi_j * \pi$ and $\pi \leq \theta_i$, we have

$$\alpha_1 \wedge \dots \wedge (\alpha_i * \pi) \wedge \dots \wedge \alpha_n = \phi_i \vee \pi = \phi_i$$

for $i = 1, \dots, n$. Thus the equalities in (2) hold. Now (1) follows from the assumption and (2). So, $(a, b) \in (\phi_1 * \pi) \vee \dots \vee (\phi_{n+1} * \pi)$.

Let consider the existence of $i \in \{1, \dots, n+1\}$ with $(a, b) \in (\phi_i * \pi)$. Then also $(u_i, v_i) \in \phi_i \cap (\phi_i * \pi)$, so we get $(u_i, v_i) \in \pi$, a contradiction. Thus for every $i \in \{1, \dots, n+1\}$ there is a nontrivial proper subquotient $r_i/s_i \subset a/b$, where $(r_i, a) \notin \pi$ or $(s_i, b) \notin \pi$ and $(r_i, s_i) \notin \pi$ such that $(r_i, s_i) \notin (\phi_i * \pi)$. Then for every $i \in \{1, \dots, n+1\}$ there is a quotient z'_i/t'_i with $r_i/s_i \rightarrow z'_i/t'_i$ and $(z'_i, t'_i) \in \phi_i$. Thus for every $i \in \{1, \dots, n+1\}$ there is a proper subquotient $r_i/s_i \subset a/b$, where $(r_i, s_i) \notin \pi$ and a quotient z_i/t_i , $(z_i, t_i) \notin \pi$ such that $r_i/s_i \rightarrow z_i/t_i$ and $u_i/v_i \rightarrow z_i/t_i$. Hence, L is $(\pi - n)$ -weakly modular.

Now, let $\pi \in \text{Con } L$, $\theta_1, \dots, \theta_n$, $\theta_i \geq \pi$ for $i = 1, \dots, n$ and $b < a$. Since

$$(a, b) \in ((\theta_1 \wedge \dots \wedge \theta_n) * \pi) \vee ((\theta_1 * \pi \wedge \dots \wedge \theta_n) * \pi) \vee \dots \vee ((\theta_1 \wedge \dots \wedge \theta_n * \pi) * \pi),$$

there is a chain $b = z_0 \leq \dots \leq z_m = a$ such that for all $i = 1, \dots, m-1$

$$(z_{i+1}, z_i) \in ((\theta_1 \wedge \dots \wedge \theta_n) * \pi) \text{ or}$$

$$(z_{i+1}, z_i) \in (\theta_1 \wedge \dots \wedge (\theta_j * \pi) \wedge \dots \wedge \theta_n) * \pi \text{ for some } j \in \{1, \dots, m\}.$$

In the first case we get (i) from the definition of $(\pi - n)$ -separability.

We should show that in the other case the condition (ii) from the definition 4.2 holds. Let $(z_{i+1}, z_i) \in (\theta_1 \wedge \dots \wedge (\theta_j * \pi) \wedge \dots \wedge \theta_n)$ for some $j \in \{1, \dots, m\}$. Further let $r/s \subset z_{i+1}/z_i$ be a nontrivial proper subquotient, $(r, s) \notin \pi$ and $(r, z_{i+1}) \notin \pi$ or $(s, z_i) \notin \pi$, and let $r/s \rightarrow u/v$, $(u, v) \notin \pi$, $(u, v) \in (\theta_1 \wedge \dots \wedge \theta_{j-1} \wedge \theta_{j+1} \wedge \dots \wedge \theta_n)$.

Suppose that for any $u' \geq v'$, the conditions $u/v \rightarrow u'/v'$ and $(u', v') \in \theta_j$ imply $(u', v') \in \pi$. By Lemma 2.2 we obtain $(u, v) \in (\theta_j * \pi)$, hence we get $(u, v) \in (\theta_1 \wedge \dots \wedge \theta_{j-1} \wedge \theta_j * \pi \wedge \theta_{j+1} \wedge \dots \wedge \theta_n)$. Since we also have $(u, v) \in ((\theta_1 \wedge \dots \wedge \theta_{j-1} \wedge \theta_j * \pi \wedge \theta_{j+1} \wedge \dots \wedge \theta_n) * \pi)$, we get $(u, v) \in \pi$, a contradiction. Therefore there exist elements $u' > v'$ such that $u/v \rightarrow u'/v'$ and $(u', v') \in \theta_j$. This yields that every (unordered) n -tuple $\theta_1, \dots, \theta_n \in \text{Con } L$, $\theta_i \geq \pi$, is $(\pi - n)$ -separable.

Conversely, let L be $(\pi - n)$ -weakly modular lattice and let every n -tuple $\theta_1, \dots, \theta_n \in \text{Con } L$, $\theta_i \geq \pi$, be $(\pi - n)$ -separable. To prove that

L satisfies the identity (L_n) it is sufficient to show that for any $b < a$
 $(a, b) \in ((\theta_1 \wedge \dots \wedge \theta_n) * \pi \vee (\theta_1 * \pi \wedge \dots \wedge \theta_n) * \pi \vee \dots \vee (\theta_1 \wedge \dots \wedge \theta_n * \pi) * \pi)$.

Let $b < a$. By $(\pi - n)$ -weakly separability of $\theta_1, \dots, \theta_n; \theta_i \geq \pi$; there exists a chain $b = c_0 \leq \dots \leq c_m = a$ such that for all $i = 0, \dots, m - 1$ either the condition (i) or the condition (ii) from the definition 4.2 holds.

In the first case we immediately obtain $(c_{i+1}, c_i) \in ((\theta_1 \wedge \dots \wedge \theta_n) * \pi)$. Now assume that (i) does not hold, so there is a quotient $u_{n+1}/v_{n+1}, (u_{n+1}, v_{n+1}) \notin \pi$ such that $c_{i+1}/c_i \rightarrow u_{n+1}/v_{n+1}$ and also $(u_{n+1}, v_{n+1}) \in (\theta_1 \wedge \dots \wedge \theta_n)$ and the condition (ii) holds. Two cases can occur:

I. there exists $j \in \{1, \dots, n\}$ such that the conditions $c_{i+1}/c_i \rightarrow u/v$ and $(u, v) \in (\theta_1 \wedge \dots \wedge (\theta_j * \pi) \wedge \dots \wedge \theta_n)$ imply $(u, v) \in \pi$. By Lemma 2.2 we get $(c_{i+1}, c_i) \in ((\theta_1 \wedge \dots \wedge \theta_j * \pi \wedge \dots \wedge \theta_n) * \pi)$

II. for every $j \in \{1, \dots, n\}$ there exists a nontrivial quotient u_j/v_j such that $c_{i+1}/c_i \rightarrow u_j/v_j$ and $(u_j, v_j) \in (\theta_1 \wedge \dots \wedge (\theta_j * \pi) \wedge \dots \wedge \theta_n)$ with $(u_j, v_j) \notin \pi$. By assumptions (i) or (ii) from the Definition 4.1 holds for the quotients $c_{i+1}/c_i, u_j/v_j, j = 1, \dots, n + 1$. The condition (i) is not satisfied, so the condition (ii) holds. Thus for every $j \in \{1, \dots, n + 1\}$ there exists a proper subquotient $r_j/s_j \subset c_{i+1}/c_i, (r_j, s_j) \notin \pi$ and $(r_j, c_{i+1}) \notin \pi$ or $(s_j, c_i) \notin \pi$, and a quotient $z_j/t_j, (z_j, t_j) \notin \pi$ such that $r_j/s_j \rightarrow z_j/t_j$ and $u_j/v_j \rightarrow z_j/t_j$. Thus for all $j = 1, \dots, n$ we get $(z_j, t_j) \in (\theta_1 \wedge \dots \wedge (\theta_j * \pi) \wedge \dots \wedge \theta_n)$. Since the condition (ii) from the Definition 4.2 holds, it follows that for some $j \in \{1, \dots, n\}$ there exists a quotient $z/t, (z, t) \notin \pi$ with $z_j/t_j \rightarrow z/t$ and $(z, t) \in \theta_j$. Since $(z_j, t_j) \in (\theta_j * \pi)$, we get $(z, t) \in (\theta_j \wedge (\theta_j * \pi))$, so $(z, t) \in \pi$, a contradiction. Therefore the case II. is impossible.

So for every $i \in \{1, \dots, m - 1\}$

$$(c_{i+1}, c_i) \in ((\theta_1 \wedge \dots \wedge \theta_n) * \pi) \text{ or}$$

$$(c_{i+1}, c_i) \in ((\theta_1 \wedge \dots \wedge \theta_j * \pi \wedge \dots \wedge \theta_n) * \pi) \text{ for some } j \in \{1, \dots, n\},$$

which yields

$$(a, b) \in ((\theta_1 \wedge \dots \wedge \theta_n) * \pi) \vee ((\theta_1 * \pi \wedge \dots \wedge \theta_n) * \pi) \vee \dots \vee ((\theta_1 \wedge \dots \wedge \theta_n * \pi) * \pi),$$

so the lattice L satisfies the identity (L'_n) . \square

As corollary we obtain the following result.

Corollary 4.4. ([6], Theorem 1) *Let L be a lattice and $n \geq 1$. Con L is (L_n) -lattice if and only if the following conditions hold:*

(i) L is $(\Delta - n)$ -weakly modular and

(ii) every n -tuple $\theta_1, \dots, \theta_n$ from $\text{Con } L$ is $(\Delta - n)$ -separable.

5. CONCLUSION

The presented work is related to the problems III.5 and III.6 of G. Grätzer's monograph [2] which ask for a characterization of lattices with Stone and (L_n) -congruence lattices for arbitrary $n \geq 1$. We present a description of arbitrary lattices whose congruence lattices considered as Heyting algebras are relative Stone (Section 3) and relative (L_n) -lattices for arbitrary $n \geq 1$ (Section 4).

We use the method of description in terms of weak projectivity of quotients introduced by G. Grätzer and E. T. Schmidt [3] and later developed by T. Katriňák and M. Haviar in [8], [4], [7], [6]. However, our method is slightly alternative to the ones presented before as it considers the congruence lattices of lattices as Heyting algebras and uses entirely the identity (L'_n) in terms of relative pseudocomplement. We present self-contained proofs for the characterizations of lattices with relative (L_n) - and relative Stone congruence lattices. As corollaries we give descriptions of lattices with Stone and (L_n) -congruence lattices for arbitrary $n \geq 1$.

REFERENCES

1. P. Crawley, Lattices whose congruence lattices form a Boolean algebra, *Pacif. J. Math.*, **10** (1960), 787–795.
2. G. Grätzer, General Lattice Theory, *Birkhäuser Verlag*, Basel 1978.
3. G. Grätzer and E. T. Schmidt, Ideals and congruence relations in lattices, *Acta Math. Acad. Sci. Hungar.*, **9** (1958), 137–175.
4. M. Haviar and T. Katriňák, Lattices whose congruence lattices is relative Stone, *Acta Sci. Math.*, (Szeged) **51** (1987), 81–91.
5. M. Haviar, Congruence lattices of lattices, Diploma work, (1988).
6. M. Haviar, Lattices whose congruence lattices satisfy Lee’s identities, *Demonstratio Math.*, **24** (1991), 247–261.
7. M. Haviar and T. Katriňák, Semilattices with (L_n) -congruence lattices, *Contribution to General Algebra 7*, (1991), 189–195.
8. T. Katriňák, Notes on Stone lattices II, *Mat. časop.*, **17** (1967), 20–37. (Russian)
9. T. Katriňák and S. El-Assar, Algebras with Boolean and Stonean congruence lattices, *Acta Math. Hungar.*, **48** (1986), 301–316.
10. K. B. Lee, Equational classes of distributive psuedocomplemented lattices, *Can. J. Math.*, **22** (1970), 881–891.
11. T. Tanaka, Canonical subdirect factorization of lattice, *J. Sci. Hiroshima Univ.*, **16** (1952), 239–246.